ELSEVIER

International Journal of Mass Spectrometry 217 (2002) 273

International Journal of Mass Spectrometry

www.elsevier.com/locate/ijms

Author index Volume 217

Alcamí, M., 119
Audier, H.E., 245

Barbieux-Flammang, M., 45, 65
Berruyer-Penaud, F., 195
Bouchoux, G., 195
Bourcier, S., 231
Bowen, R.D., 109
Burgers, P.C., 97

Cao, J., 179
Choret, N., 195
Correia, A.J.F., 55

Decouzon, M., 75
Denhez, J.P., 245

Fernandes, A.M., 55
Flammang, R., 1, 45, 65, 195
Fokkens, R.H., 55
Fura, A., 81
Fürst, T., 153

Gal, J.-F., 75
Gerbaux, P., 1, 45, 65
Griffin, L.L., 23
Grüzmacher, H.-F., 153
Gualano, E., 65

Harrison, A.G., 185
Heydorn, L.N., 109
Holmes, J.L., 179
Hoppilliard, Y., 231
Hudson, C.E., 23

Kong, J., 257
Kuck, D., 131

Lavorato, D.J., 1
Le, H.T., 45, 65
Leblanc, D., 245
Letzel, M.C., 153
Loos, J., 169
Luna, A., 119

Maria, P.-C., 75
Matthias, C., 131
Mayer, K.K., 153
Mayer, P.S., 257
McAdoo, D.J., 23
McLafferty, F.W., 81
Mó, O., 119
Morton, T.H., 257

Nedev, H., 245
Nguyen, M.T., 45, 65
Nibbering, N.M.M., 55

Ruttink, P.J.A., 97

Schröder, D., 169
Schwarz, H., 169
Sun, W., 179

Terlouw, J.K., 1, 97, 109
Tortajada, J., 119
Traeger, J.C., 23
Trikoupis, M.A., 1, 97
Tureček, F., 65, 81

Wiegrebe, W., 153

Yáñez, M., 119

Zummack, W., 169

PII S1387-3806(02)00677-2

ELSEVIER

International Journal of Mass Spectrometry 217 (2002) 275–277

International Journal of Mass Spectrometry

www.elsevier.com/locate/ijms

Subject index Volume 217

Ab initio
Functional group migration in benzaldoxime-*O*-*n*-propyl ether radical cation, 153
Ab initio calculations
Self-catalysis in the gas-phase: enolization of the acetone radical cation, 97
Acid-catalyzed cyclization
Thermochemistry and unimolecular reactivity of protonated α,ω-aminoalcohols in the gas phase, 195
Adduct
Adduct formation between phthalate esters and Li^+ in the gas phase: a thermochemical study by FT-ICR mass spectrometry, 75
O-alkyl benzaldoximes
Functional group migration in benzaldoxime-*O*-*n*-propyl ether radical cation, 153
Alkyl transfer transition states
Carbocation rearrangements of trimethylsilyl adducts of saturated acyclic C_5–C_7 ketones in the gas phase, 257
Allylic cations
Carbocation rearrangements of trimethylsilyl adducts of saturated acyclic C_5–C_7 ketones in the gas phase, 257
Ammonia loss
Evidence for the formation of acyclic ions from the radical cations and cyclic ions from the protonated molecules of α,ω-diamines upon loss of ammonia, 55
Aniline
Ionized aniline and its distonic radical cation isomers, 45
Associative ion–molecule reactions
Hydrogen-shift isomers of ionic and neutral hydroxypyridines: a combined experimental and computational investigation, 1

Basicity
Adduct formation between phthalate esters and Li^+ in the gas phase: a thermochemical study by FT-ICR mass spectrometry, 75
Benzaldehyde
Distonic isomers of ionized benzaldehyde, 65
Binding energies
The ions $CH_6O^{\bullet+}$ and CH_7O^+, 179

Carbon
Silicon- versus carbon-containing ions: 1,3-CH_3^+ transfers, 245
Carbon cluster
Small carbon clusters (C_n^0, C_n^+, C_n^-) from acyclic and cyclic precursors. Neutralization–reionization and theory, 81
C–H bond activation
Regio- and diastereoselective C–H bond activation of valeramide and 3-methyl valeramide by bare Fe^+ ions, 169
CH_3^+ transfers
Silicon- versus carbon-containing ions: 1,3-CH_3^+ transfers, 245
Chemical ionization
Carbocation rearrangements of trimethylsilyl adducts of saturated acyclic C_5–C_7 ketones in the gas phase, 257
Collisionally activated decomposition
Carbocation rearrangements of trimethylsilyl adducts of saturated acyclic C_5–C_7 ketones in the gas phase, 257
Collisionally activated dissociation
Small carbon clusters (C_n^0, C_n^+, C_n^-) from acyclic and cyclic precursors. Neutralization–reionization and theory, 81
Collision-induced dissociation
Evidence for the formation of acyclic ions from the radical cations and cyclic ions from the protonated molecules of α,ω-diamines upon loss of ammonia, 55
Reactions of low-energy pentenyl methyl ether radical cations $C_2H_5CH{=}CHCH_2OCH_3^{\bullet+}$, $CH_2{=}CHCH(C_2H_5)OCH_3^{\bullet+}$ and $CH_2{=}C(C_2H_5)CH_2OCH_3^{\bullet+}$, 109
Convergency
A theoretical study of the interaction between Ni^+ and small oxygen- and nitrogen-containing bases, 119
Cyclization reactions
Large hydrocarbon ion/molecule complexes formed during the unimolecular fragmentation of protonated *tert*-butyl-substituted tri- and tetrabenzylmethane, 131
Cyclopropylcarbinyl cations
Carbocation rearrangements of trimethylsilyl adducts of saturated acyclic C_5–C_7 ketones in the gas phase, 257

Density functional theory
Why are smaller fragments preferentially lost from radical cations at low energies and larger ones at high energies?. An experimental and theoretical study, 23
A theoretical study of the interaction between Ni^+ and small oxygen- and nitrogen-containing bases, 119
Density-functional calculations
Hydrogen-shift isomers of ionic and neutral hydroxypyridines: a combined experimental and computational investigation, 1

PII S1387-3806(02)00678-4

Deuterated derivatives
Functional group migration in benzaldoxime-*O*-*n*-propyl ether radical cation, 153
Deuterium labelling
Reactions of low-energy pentenyl methyl ether radical cations $C_2H_5CH{=}CHCH_2OCH_3^{\bullet+}$, $CH_2{=}CHCH(C_2H_5)OCH_3^{\bullet+}$ and $CH_2{=}C(C_2H_5)CH_2OCH_3^{\bullet+}$, 109
DFT molecular orbital calculations
B3LYP DFT molecular orbital approach, an efficient method to evaluate the thermochemical properties of MALDI matrices, 231
α,ω-Diamines
Evidence for the formation of acyclic ions from the radical cations and cyclic ions from the protonated molecules of α,ω-diamines upon loss of ammonia, 55
Dimer radical cation
Self-catalysis in the gas-phase: enolization of the acetone radical cation, 97
Distonic ion
Functional group migration in benzaldoxime-*O*-*n*-propyl ether radical cation, 153
Distonic ions
Hydrogen-shift isomers of ionic and neutral hydroxypyridines: a combined experimental and computational investigation, 1
Distonic isomers of ionized benzaldehyde, 65
Reactions of low-energy pentenyl methyl ether radical cations $C_2H_5CH{=}CHCH_2OCH_3^{\bullet+}$, $CH_2{=}CHCH(C_2H_5)OCH_3^{\bullet+}$ and $CH_2{=}C(C_2H_5)CH_2OCH_3^{\bullet+}$, 109
Distonic radical cations
Ionized aniline and its distonic radical cation isomers, 45

Electron affinity
B3LYP DFT molecular orbital approach, an efficient method to evaluate the thermochemical properties of MALDI matrices, 231
Enolization
Self-catalysis in the gas-phase: enolization of the acetone radical cation, 97

Fe^+ chemistry
Regio- and diastereoselective C–H bond activation of valeramide and 3-methyl valeramide by bare Fe^+ ions, 169
Fragmentation mechanism
Functional group migration in benzaldoxime-*O*-*n*-propyl ether radical cation, 153
Fragmentation reactions
Fragmentation reactions of protonated peptides containing phenylalanine: a linear free energy correlation in the fragmentation of H–Gly–Xxx–Phe–OH, 185
FT-ICR
Adduct formation between phthalate esters and Li^+ in the gas phase: a thermochemical study by FT-ICR mass spectrometry, 75

H/D exchange
Large hydrocarbon ion/molecule complexes formed during the unimolecular fragmentation of protonated *tert*-butyl-substituted tri- and tetrabenzylmethane, 131
Hydride transfer
Large hydrocarbon ion/molecule complexes formed during the unimolecular fragmentation of protonated *tert*-butyl-substituted tri- and tetrabenzylmethane, 131
Hydride transfer transition states
Carbocation rearrangements of trimethylsilyl adducts of saturated acyclic C_5–C_7 ketones in the gas phase, 257
Hydroxypyridine
Hydrogen-shift isomers of ionic and neutral hydroxypyridines: a combined experimental and computational investigation, 1

Ion structures
Evidence for the formation of acyclic ions from the radical cations and cyclic ions from the protonated molecules of α,ω-diamines upon loss of ammonia, 55
Ion/molecule complexes
Large hydrocarbon ion/molecule complexes formed during the unimolecular fragmentation of protonated *tert*-butyl-substituted tri- and tetrabenzylmethane, 131
Ion/molecule reactions
Evidence for the formation of acyclic ions from the radical cations and cyclic ions from the protonated molecules of α,ω-diamines upon loss of ammonia, 55
Ion/neutral complexes
The ions $CH_6O^{\bullet+}$ and CH_7O^+, 179
Ionization energy
B3LYP DFT molecular orbital approach, an efficient method to evaluate the thermochemical properties of MALDI matrices, 231

Lithium cation
Adduct formation between phthalate esters and Li^+ in the gas phase: a thermochemical study by FT-ICR mass spectrometry, 75

MALDI matrices
B3LYP DFT molecular orbital approach, an efficient method to evaluate the thermochemical properties of MALDI matrices, 231
Mass spectrometry
B3LYP DFT molecular orbital approach, an efficient method to evaluate the thermochemical properties of MALDI matrices, 231
Metastable ion
Functional group migration in benzaldoxime-*O*-*n*-propyl ether radical cation, 153
Metastable ion decomposition
Carbocation rearrangements of trimethylsilyl adducts of saturated acyclic C_5–C_7 ketones in the gas phase, 257
Metastable ions
Large hydrocarbon ion/molecule complexes formed during the unimolecular fragmentation of protonated *tert*-butyl-substituted tri- and tetrabenzylmethane, 131
MIKE spectra
Carbocation rearrangements of trimethylsilyl adducts of saturated acyclic C_5–C_7 ketones in the gas phase, 257

Neutralisation–reionisation mass spectrometry
Hydrogen-shift isomers of ionic and neutral hydroxypyridines: a combined experimental and computational investigation, 1

Neutralization–reionization
Small carbon clusters (C_n^0, C_n^+, C_n^-) from acyclic and cyclic precursors. Neutralization–reionization and theory, 81
Ni^+ binding energies
A theoretical study of the interaction between Ni^+ and small oxygen- and nitrogen-containing bases, 119

Phenylalanine residues
Fragmentation reactions of protonated peptides containing phenylalanine: a linear free energy correlation in the fragmentation of H–Gly–Xxx–Phe–OH, 185
Photoionization mass spectrometry
Why are smaller fragments preferentially lost from radical cations at low energies and larger ones at high energies? An experimental and theoretical study, 23
Phthalate
Adduct formation between phthalate esters and Li^+ in the gas phase: a thermochemical study by FT-ICR mass spectrometry, 75
Proton affinities
Ionized aniline and its distonic radical cation isomers, 45
Proton affinity
B3LYP DFT molecular orbital approach, an efficient method to evaluate the thermochemical properties of MALDI matrices, 231
Proton transfer
Large hydrocarbon ion/molecule complexes formed during the unimolecular fragmentation of protonated *tert*-butyl-substituted tri- and tetrabenzylmethane, 131
Proton transfer catalysis
Self-catalysis in the gas-phase: enolization of the acetone radical cation, 97
Protonated α,ω-aminoalcohols
Thermochemistry and unimolecular reactivity of protonated α,ω-aminoalcohols in the gas phase, 195
Protonated ketones
Carbocation rearrangements of trimethylsilyl adducts of saturated acyclic C_5–C_7 ketones in the gas phase, 257
Protonated peptides
Fragmentation reactions of protonated peptides containing phenylalanine: a linear free energy correlation in the fragmentation of H–Gly–Xxx–Phe–OH, 185

Quantum chemical calculations
Ionized aniline and its distonic radical cation isomers, 45
Quantum mechanical calculations
Distonic isomers of ionized benzaldehyde, 65

Rearrangement
Reactions of low-energy pentenyl methyl ether radical cations $C_2H_5CH{=}CHCH_2OCH_3^{\bullet+}$, $CH_2{=}CHCH(C_2H_5)OCH_3^{\bullet+}$ and $CH_2{=}C(C_2H_5)CH_2OCH_3^{\bullet+}$, 109
Functional group migration in benzaldoxime-*O*-*n*-propyl ether radical cation, 153
RRKM theory
Why are smaller fragments preferentially lost from radical cations at low energies and larger ones at high energies? An experimental and theoretical study, 23

Silicon
Silicon- versus carbon-containing ions: 1,3-CH_3^+ transfers, 245
Size effects
Why are smaller fragments preferentially lost from radical cations at low energies and larger ones at high energies? An experimental and theoretical study, 23
Spin contamination
A theoretical study of the interaction between Ni^+ and small oxygen- and nitrogen-containing bases, 119
Stereochemistry
Regio- and diastereoselective C–H bond activation of valeramide and 3-methyl valeramide by bare Fe^+ ions, 169
Structures
The ions $CH_6O^{\bullet+}$ and CH_7O^+, 179
Surface crossing
Self-catalysis in the gas-phase: enolization of the acetone radical cation, 97

Tandem mass spectrometer
Ionized aniline and its distonic radical cation isomers, 45
Tandem mass spectrometry
Distonic isomers of ionized benzaldehyde, 65
Self-catalysis in the gas-phase: enolization of the acetone radical cation, 97
tert-Butylbenzenes
Large hydrocarbon ion/molecule complexes formed during the unimolecular fragmentation of protonated *tert*-butyl-substituted tri- and tetrabenzylmethane, 131
Transition states
Why are smaller fragments preferentially lost from radical cations at low energies and larger ones at high energies? An experimental and theoretical study, 23